

# Biosolids Calculations



Rachel Barnes-McAden



Solids	30.94	309400
Nitrogen (TKN)	4.45	44500
Phosphorus	1.72	17200
Potassium	0.20	2000
Sulfur	0.60	6000
Calcium	9.86	98600
Magnesium	0.29	2900
Sodium	0.10	1000
Iron		49600
Manganese		178
Copper		269
Zinc		421
Ammonia Nitrogen	0.27	2700
NO <sub>3</sub> -NO <sub>2</sub> Nitrogen		21
Cadmium		2.0
Chromium		49
Nickel		19
Lead		40
Arsenic		2.15
Mercury		0.96
Selenium		2.23
pH (Standard Units)	12.10	
Calcium Carbonate Eq	14.63	146300
Volatile Solids	64.88	648800
Organic Nitrogen	4.18	41800
Molybdenum		13

All Values, except for Solids, are on a Dry Weight Basis.



The biosolids analysis reports nutrient levels in terms of the percent by weight.

We're going to figure out how much of each nutrient there is in terms of pounds per dry ton.

For % to decimal : divide by 100

(or use % button on calculator)



## Calculating Nitrogen

$$0.045 \text{ (TKN)} \times 2,000 \text{ lbs./T} = 89.0 \text{ lbs TKN/DT}$$

(4.45%)

$$0.0027 \text{ (NH}_3\text{)} \times 2,000 \text{ lbs./T} = 5.4 \text{ lbs NH}_3\text{/DT}$$

(0.27%)

$$89.0 \text{ TKN/DT} - 5.4 \text{ lbs NH}_3\text{/DT} = 83.6 \text{ lbs organic N/DT}$$

# Nitrogen Calculations for Ammonium, Organic and Residual Nitrogen Based on Analysis of Material

Material: Biosolids

Analysis:  TKN  NH<sub>4</sub>-N  P<sub>2</sub>O<sub>5</sub>  K<sub>2</sub>O \ ton or 1,000 gals. DT

Days before

Incorporation: (circle one)

Injected, Broadcast-Immediate Incorp., >2, >4, >7 or No Incorp., Irrigate-No Incorp.

Previous Application: 0-1 yr. of the last 5, 2-3 yrs. of the last 5, 4-5 yrs. of the last 5

## First Year - Plant Available Nitrogen (PAN)

<div style="border: 1px solid black; display: inline-block; padding: 2px 10px;">89</div> DT TKN/unit
<div style="border: 1px solid black; display: inline-block; padding: 2px 10px;">5.4</div> DT - NH <sub>4</sub> -N/unit
<div style="border: 1px solid black; display: inline-block; padding: 2px 10px;">83.6</div> DT Organic N/unit

<div style="border: 1px solid black; display: inline-block; width: 60px; height: 25px; margin: 0 auto;"></div> = x availability coefficient	<div style="border: 1px solid black; display: inline-block; width: 100px; height: 25px; margin: 0 auto;"></div> #/unit
<div style="border: 1px solid black; display: inline-block; width: 60px; height: 25px; margin: 0 auto;"></div> = x availability coefficient	<div style="border: 1px solid black; display: inline-block; width: 100px; height: 25px; margin: 0 auto;"></div> #/unit
<div style="border: 1px solid black; display: inline-block; width: 150px; height: 25px; margin: 0 auto;"></div> TOTAL PAN	<div style="border: 1px solid black; display: inline-block; width: 100px; height: 25px; margin: 0 auto;"></div> #/unit

Availability Coefficients from Standards and Criteria

Manure	Biosolids
Table 8-2	Table 9-2
Table 8-2	Table 9-1

           #/unit x            units/ac. =            # PAN/ac.

Unit = Ton or 1,000 Gallons

**Table 9-1**  
**Estimated Nitrogen Mineralization Rates for Biosolids<sup>1</sup>**  
 (S&C pg 117)

Biosolids Type	Application Year			
	Application Year	1 Year After Application	2 Years After Application	3 Years After Application
Lime Stabilized	0.30 	0.10	0.10	0.05
Aerobic Digestion	0.30	0.10	0.10	0.05
Anaerobic Digestion	0.30	0.10	0.10	0.05
Composted <sup>2</sup>	0.10	0.05	0.03	0.00

**Table 9-2**  
**Biosolids Ammonium Nitrogen Availability Coefficients<sup>1</sup>**  
 (S&C pg 117)

Method of Application	Biosolids pH < 10	Biosolids pH > 10
Injection	1.00	1.00
Incorporated within 24 hours	0.85	0.75 
Incorporated within 1-7 days	0.70	0.50
Incorporated after 7 days or no incorporation	0.50	0.25

## First Year - Plant Available Nitrogen (**PAN**)

89 DT
TKN/unit
- 5.4 DT
NH <sub>4</sub> -N/unit
83.6 DT
Organic N/unit

0.75 =  
x availability coefficient

\_\_\_\_ #/unit

0.30 =  
x availability coefficient

\_\_\_\_ #/unit

TOTAL PAN

\_\_\_\_ #/unit

### Availability Coefficients from Standards and Criteria

Manure	Biosolids
--------	-----------

Table 8-2	Table 9-2
-----------	-----------

Table 8-2	Table 9-1
-----------	-----------

\_\_\_\_ #/unit x \_\_\_\_\_ units/ac. = \_\_\_\_\_ # PAN/ac.

Unit = Ton or 1,000 Gallons

# First Year - Plant Available Nitrogen (**PAN**)

<b>89</b> DT
TKN/unit
- <b>5.4</b> DT
NH <sub>4</sub> -N/unit
<b>83.6</b> DT
Organic N/unit

<b>0.75</b>	=	<b>4.05</b> #/unit
x availability coefficient		
<b>0.30</b>	=	<b>25.08</b> #/unit
x availability coefficient		
<b>TOTAL PAN</b>		<b>29.13</b> #/unit

## Availability Coefficients from Standards and Criteria

<u>Manure</u>	<u>Biosolids</u>
Table 8-2	Table 9-2
Table 8-2	Table 9-1



## NUTRIENT MANAGEMENT Balance Sheet

Name: WillDaLyn Farms  
Tract: T-1989

Date: February 2010

Field Name	Ac.	Crop Rotation	Expctd Yield (bu or tons)	Nutrient Needs (from soil test & expctd yield) N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O	Nitrogen Residual (Leg./ Manure)	Days before Incorp	Organic Material Applied (1000 gal. or tons/ac)	Org. Nut. Applied N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O	N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O Need or (Surplus)	N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O (commercial)	Notes
HF-1	8	Orchard-grass Pasture	3.6 ac/au	50-40-70	0/14	----	----	----	36-40-70	36-40- 70 br	
HF-2A	16	Corn (grain)	120 bu/ac.	120-80-100	0/14	>7					
HF-2B	12	Corn (grain)	90 bu/ac	90-0-0	0/0	>2					
HF-2C	9	Orchard-grass Hay (maint.)	3.3 t/ac	140-40-95	0/0	>7					1
HF-3A	11	Corn (grain)	150 bu/ac	150-120-100	0/14	>1					
HF-3B	11.3	Corn (silage)	15.6 t/ac	90-100-200	0/7	>2	6 k/ac Dairy	63-69-119	20-31-81	0-0-81 br 20-31-0 ba	

**Notes:** Application Methods: Ba= Banded, Br = Broadcast, Sd = Sidedress, k = 1000 gal.

1. Nitrogen needs are shown for 2 cuttings of hay.



## Biosolids Application Rate

From case study, **Field 3A** is a **corn** (grain) field.

Crop nutrient need is **150-120-100**.

First, credit residual nitrogen carryover to crop nutrient needs:

$$150 \text{ lbs N} - 14 \text{ lbs N} = \mathbf{136 \text{ lbs N need}}$$

(dairy manure residual)

Balance of crop nutrient need is **136-120-100**

## How Many Tons of Biosolids to Apply ?

Field Name	Acres	Crop Rotation	Expected Yield Bu/ Ton	Nutrient Needs (soil test) N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O	Nitrogen Residual (Leg./ Manure)	Days before Incorp	Material Applied (1,000 gal. or Tons/ac)	Org. Nut. Applied N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O	N N (S
HF-2C	9	Orchard grass Hay	3.3 t/ac	140-40-95	0/0	>7			
HF-3A	11	Corn (grain)	150 bu/ac	150-120-100	0/14	>1			
HF-3B	11.3	Corn (silage)	15.6 t/ac	90-100-200	0/7	>2	6k Dairy	63-69-119	

$$150 \text{ lbs N for corn} - 14 \text{ lbs N residual} = 136 \text{ lbs N from Biosolids}$$



**Nitrogen** needs are to be met through biosolids application

From worksheet, we get **29.13 lbs PAN/ DT**

Crop needs, less residual, are **136 lbs N**

(14 lbs. dairy manure)

$$\frac{136 \text{ lbs N/ Acre}}{29.13 \text{ lbs N/ DT}} = 4.67 \text{ DT/ A}$$


# Fill in Worksheet


$$\frac{136}{\#N/ac. \text{ needed}} \div \frac{29.13}{PAN\#/DT} \frac{1,000 \text{ gallons or tons}}{\text{---}} = \frac{4.67}{\text{---}} \frac{1,000 \text{ gallons or tons}}{\text{---} \text{ acre}}$$

Unit = Ton or 1,000 Gallons



Remember that the biosolids will arrive as **wet tons**, but we just calculated dry tons. So- we'll need to convert:

$$\frac{\text{DT}}{\% \text{ solids}} = \text{WT}$$

$$\frac{4.67 \text{ DT/ A}}{0.3094} = 15.09 \text{ WT/ A}$$

(30.94% solids)

↑  
to achieve **N = 136 lbs/ A**

# NUTRIENT MANAGEMENT Balance Sheet

Name: WillDaLyn Farms  
 Address: T-1989

Date: February 2010

Field Name	Ac.	Crop Rotation	Expctd Yield (bu or tons)	Nutrient Needs (from soil test & expctd yield) N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O	Nitrogen Residual (Leg./Manure)	Days before Incorporation	Organic Material Applied (1000 gal. or tons/ac)	Org. Nut. Applied N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O	N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O Need or (Surplus)	N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O (commercial)	Notes
-1	8	Orchard-grass Pasture	3.6 ac/au	50-40-70	0/14	---	---	---	36-40-70	36-40-70 br	
-2A	16	Corn (grain)	120 bu/ac.	120-80-100	0/14	>7					
-2B	12	Corn (grain)	90 bu/ac	90-0-0	0/0	>2					
-2C	9	Orchard-grass Hay (maint.)	3.3 t/ac	140-40-95	0/0	>7					1
-3A	11	Corn (grain)	150 bu/ac	150-120-100	0/14	>1	4.67 DT 15.09 WT/ac	136-?-?			
-3B	11.3	Corn (silage)	15.6 t/ac	90-100-200	0/7	>2	6 k/ac Dairy	63-69-119	20-31-81	0-0-81 br 20-31-0 ba	

Notes: Application Methods: Ba= Banded, Br = Broadcast, Sd = Sidedress, k = 1000 gal.

1. Nitrogen needs are shown for 2 cuttings of hay.



## Calculating Phosphorus

1.72 % elemental P

$$0.0172 \text{ P} \times 2,000 \text{ lbs/ T} = 34.4 \text{ lbs P/ DT}$$

For field nutrients, P is dealt with as  $\text{P}_2\text{O}_5$  (phosphate)

$$\text{P} \times 2.29 = \text{P}_2\text{O}_5$$

(conversion factor)

$$34.4 \text{ lbs P/ DT} \times 2.29 = \mathbf{78.8 \text{ lbs P}_2\text{O}_5/ \text{DT}}$$





## Calculating Potassium

0.20 % elemental K

$$0.0020 \text{ K} \times 2,000 \text{ lbs/ T} = 4.0 \text{ lbs K/ DT}$$

For field nutrients, K is dealt with as  $\text{K}_2\text{O}$  (Potash)

$$\text{K} \times \underset{\text{(conversion factor)}}{1.2} = \text{K}_2\text{O}$$

$$4.0 \text{ lbs K/ DT} \times 1.2 = \mathbf{4.8 \text{ lbs K}_2\text{O/ DT}}$$




We'll need to know how much

**Phosphate** and **Potash** will be applied when biosolids are used to meet the **Nitrogen** needs of the corn.

$$78.8 \text{ lbs P}_2\text{O}_5/\text{DT} \times 4.67 \text{ DT/A} = \mathbf{368 \text{ lbs P}_2\text{O}_5/\text{A}}$$


$$4.8 \text{ lbs K}_2\text{O}/\text{DT} \times 4.67 \text{ DT/A} = \mathbf{22 \text{ lbs K}_2\text{O}/\text{A}}$$




Original Crop Needs: 150-120-100

<u>Nutrient</u>	<u>Amount/ Source</u>	<u>Net</u>
<b>Nitrogen:</b> 150	14 lbs/ A from Residual	136
	136 lbs/ A from Biosolids	0
<b>Phosphate:</b> 120	368 lbs/A from Biosolids	+248*
<b>Potash:</b> 100	22 lbs/ A from Biosolids	- 78
	78 lbs/ A from fertilizer	0

\* Can be “banked” for crops in remainder of rotation (3 yrs X 120 #P<sub>2</sub>O<sub>5</sub>/yr)

# NUTRIENT MANAGEMENT Balance Sheet

Name: WillDaLyn Farms  
 Address: T-1989

Date: February 2010

Field Name	Ac.	Crop Rotation	Expctd Yield (bu or tons)	Nutrient Needs (from soil test & expctd yield) N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O	Nitrogen Residual (Leg./Manure)	Days before Incorporation	Organic Material Applied (1000 gal. or tons/ac)	Org. Nut. Applied N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O	N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O Need or (Surplus)	N-P <sub>2</sub> O <sub>5</sub> -K <sub>2</sub> O (commercial)	Notes
-1	8	Orchard-grass Pasture	3.6 ac/au	50-40-70	0/14	---	---	---	36-40-70	36-40-70 br	
-2A	16	Corn (grain)	120 bu/ac.	120-80-100	0/14	>7					
-2B	12	Corn (grain)	90 bu/ac	90-0-0	0/0	>2					
-2C	9	Orchard-grass Hay (maint.)	3.3 t/ac	140-40-95	0/0	>7					1
-3A	11	Corn (grain)	150 bu/ac	150-120-100	0/14	>1	4.67 DT 15.09 WT/ac	136-368-22	0-(248*)-78	0-0-78	*P
-3B	11.3	Corn (silage)	15.6 t/ac	90-100-200	0/7	>2	6 k/ac Dairy	63-69-119	20-31-81	0-0-81 br 20-31-0 ba	

Notes: Application Methods: Ba= Banded, Br = Broadcast, Sd = Sidedress, k = 1000 gal.

1. Nitrogen needs are shown for 2 cuttings of hay.

\*P-Ensure this amount can be used during this crop rotation.



# Lime Applied

- Calculating Lime Application

14.63 % Calcium Carbonate Equiv.

(from biosolids analysis)

4.67 DT/A X .1463 = **.68 T/A** of Lime

# Biosolids Residual

## Nitrogen Calculations for Ammonium, Organic and Residual Nitrogen Based on Analysis of Material

Material: Biosolids

Analysis:  TKN  NH<sub>4</sub>-N  P.C.  K.O. <sup>DT</sup> \ ton or 1,000 gals.

Days before

Incorporation: (circle one)

Injected, Broadcast-Immediate Incorp. >2, >4, >7 or No Incorp., Irrigate-No Incorp.

Previous Application: 0-1 yr. of the last 5, 2-3 yrs. of the last 5, 4-5 yrs. of the last 5

### First Year - Plant Available Nitrogen (**PAN**)

<b>89</b> <sup>DT</sup>
TKN/unit
- <b>5.4</b> <sup>DT</sup>
NH <sub>4</sub> -N/unit
<b>83.6</b> <sup>DT</sup>
Organic N/unit

<input type="text"/>	=	<input type="text"/> #/unit
x availability coefficient		
<input type="text"/>	=	<input type="text"/> #/unit
x availability coefficient		
<b>TOTAL PAN</b>		<input type="text"/> #/unit

Availability Coefficients from Standards and Criteria

Manure	Biosolids
Table 8-2	Table 9-2
Table 8-2	Table 9-1

           #/unit x            units/ac. =            # PAN/ac.

Unit = Ton or 1,000 Gallons

# Use Table 9.1

Ammonium = 5.4 lbs.N/ Dry Ton      TKN = 89 lbs. N/ DT

Organic N = 83.6 lbs. N/ DT

## Section IX. Biosolids Management

**Table 9-1  
Estimated Nitrogen Mineralization Rates for Biosolids<sup>1</sup>**

Biosolids Type	Application Year			
	Application Year	1 Year After Application	2 Years After Application	3 Years After Application
Lime Stabilized	0.30	0.10	0.10	0.05
Aerobic Digestion	0.30	0.10	0.10	0.05
Anaerobic Digestion	0.30	0.10	0.10	0.05
Composted <sup>2</sup>	0.10	0.05	0.03	0.00

1. To determine nitrogen available from previous Biosolids applications, multiply the percent organic nitrogen by the appropriate mineralization factor.
2. Total organic nitrogen content of 2% or less and no significant ammonia nitrogen.

**Table 9-2  
Biosolids Ammonium Nitrogen Availability Coefficients<sup>1</sup>**

Method of Application	Biosolids pH < 10	Biosolids pH > 10
Injection	1.00	1.00
Incorporated within 24 hours	0.85	0.75
Incorporated within 1-7 days	0.70	0.50
Incorporated after 7 days or no incorporation	0.50	0.25

1. To determine the plant-available Biosolids ammonium nitrogen in the soil, multiply the Biosolids ammonium nitrogen concentration or total weight applied by the appropriate availability coefficient.

# To Calculate Biosolids Residual One Year after Application

Residual - Plant Available Nitrogen (for following year)

83.6	X	0.10	=	8.36	#/ <sup>DT</sup> unit
Organic N/ <sup>DT</sup> unit	x	availability coefficient			

<b>Manure</b>	<b>Biosolids</b>
Table 8-3	Table 9-1

\_\_\_\_\_ #/unit x \_\_\_\_\_ units/ac. = \_\_\_\_\_ # Residual Nitrogen/ac.

## Completed Worksheet

Residual - Plant Available Nitrogen (for following year)

$$\boxed{83.6} \text{ Organic N/unit} \times \boxed{0.10} \text{ availability coefficient} = \underline{8.36} \text{ \# /unit}^{\text{DT}}$$

Manure	Biosolids
Table 8-3	Table 9-1

$$\underline{8.36} \text{ \# /unit}^{\text{DT}} \times \underline{4.67} \text{ units/ac.}^{\text{DT}} = \underline{39.03} \text{ \# Residual Nitrogen/ac.}$$



Questions ?